

Measuring Complexity of Maximal Matchings of Graphs

Oscar Levin

University of Northern Colorado

Computability Theory and its Applications
University of Waterloo, 2018

Joint work with Stephen Flood, Matthew Jura, and Tyler Markkanen.

Matchings in Graphs

Definition

A *matching* in a graph $G = (V, E)$ is a subset $M \subseteq E$ for which each vertex is incident to at most one edge in M .


A matching is *perfect* if every vertex is incident to exactly one edge in M .

Matchings in Graphs

Definition

A *matching* in a graph $G = (V, E)$ is a subset $M \subseteq E$ for which each vertex is incident to at most one edge in M .

A matching is *perfect* if every vertex is incident to exactly one edge in M .




Matchings in Graphs

Definition

A *matching* in a graph $G = (V, E)$ is a subset $M \subseteq E$ for which each vertex is incident to at most one edge in M .

A matching is *perfect* if every vertex is incident to exactly one edge in M .




Matchings in Graphs

Definition

A *matching* in a graph $G = (V, E)$ is a subset $M \subseteq E$ for which each vertex is incident to at most one edge in M .

A matching is *perfect* if every vertex is incident to exactly one edge in M .



Maximal matchings

Not every graph contains a perfect matching.

Maximal matchings

Not every graph contains a perfect matching.

Theorem (Steffens)

Every graph contains a maximal matching.

Maximal matchings

Not every graph contains a perfect matching.

Theorem (Steffens)

Every graph contains a maximal matching.

Definition

- ▶ A matching M is (*weakly*) *maximal* provided there is no matching N with $M \subset N$.

Maximal matchings

Not every graph contains a perfect matching.

Theorem (Steffens)

Every graph contains a maximal matching.

Definition

- ▶ A matching M is *(weakly) maximal* provided there is no matching N with $M \subset N$.
- ▶ A matching M is *maximal* provided there is no matching N with $\text{supp}(M) \subset \text{supp}(N)$.
($\text{supp}(M)$ is the set of vertices incident to M .)

Maximal matchings

Not every graph contains a perfect matching.

Theorem (Steffens)

Every graph contains a maximal matching.

Definition

- ▶ A matching M is *(weakly) maximal* provided there is no matching N with $M \subset N$.
- ▶ A matching M is *maximal* provided there is no matching N with $\text{supp}(M) \subset \text{supp}(N)$.
($\text{supp}(M)$ is the set of vertices incident to M .)


Improving matchings

How can you augment a matching?



Improving matchings

How can you augment a matching?




Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

Improving matchings

How can you augment a matching?



Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

Improving matchings

How can you augment a matching?



Definition


An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition


An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition


An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition


An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Improving matchings

How can you augment a matching?



Definition

An M -alternating path has edges alternating in and out of M .

An M -augmenting path is an M -alternating path that starts with an unmatched vertex and either ends in another unmatched vertex or is infinite.

If there is an M -augmenting path starting at v , then there is a matching M' that improves the support of M to include v (and possibly one other vertex).

Steffens' Theorem

Theorem (Steffens)

A graph has a perfect matching iff for any matching M and unmatched v , there is an M -augmenting path starting at v .

Steffens' Theorem

Theorem (Steffens)

A graph has a perfect matching iff for any matching M and unmatched v , there is an M -augmenting path starting at v .

↖ Condition (A)

Steffens' Theorem

Theorem (Steffens)

A graph has a perfect matching iff for any matching M and unmatched v , there is an M -augmenting path starting at v .

↖ Condition (A)

Corollary

Every graph has a maximal matching.

Steffens' Theorem

Theorem (Steffens)

A graph has a perfect matching iff for any matching M and unmatched v , there is an M -augmenting path starting at v .

↖ Condition (A)

Corollary

Every graph has a maximal matching.

Really, maximality seems to be a corollary to the proof of this theorem.

Our goal

Use Reverse Mathematics to understand the strength of both of these theorems.

Our goal

Use Reverse Mathematics to understand the strength of both of these theorems.

The plan:

1. Complete classification for locally finite graphs.

Our goal

Use Reverse Mathematics to understand the strength of both of these theorems.

The plan:

1. Complete classification for locally finite graphs.
2. Get a sense why the general case is much much much much harder to classify (probably).

Definition

- ▶ A graph is *locally finite* provided every vertex has finite degree.
- ▶ A graph is *bounded* provided there is a function $h : V \rightarrow \mathbb{N}$ s.t. $\forall x, y \in V(\{x, y\} \in E \rightarrow h(x) \geq y)$.

Locally finite and bounded graphs

Definition

- ▶ A graph is *locally finite* provided every vertex has finite degree.
- ▶ A graph is *bounded* provided there is a function $h : V \rightarrow \mathbb{N}$ s.t. $\forall x, y \in V(\{x, y\} \in E \rightarrow h(x) \geq y)$.

Think: bounded = highly computable.

Locally finite and bounded graphs

Theorem

The following are equivalent over RCA_0 :

- 1. Every locally finite graph has a maximal matching.*
- 2. A locally finite graph has a perfect matching iff it satisfies condition (A).*
- 3. ACA_0 .*

Locally finite and bounded graphs

Theorem

The following are equivalent over RCA_0 :

- 1. Every locally finite graph has a maximal matching.*
- 2. A locally finite graph has a perfect matching iff it satisfies condition (A).*
- 3. ACA_0 .*

Theorem

The following are equivalent over RCA_0 :

- 1. Every **bounded** graph has a maximal matching.*
- 2. A **bounded** graph has a perfect matching iff it satisfies condition (A).*
- 3. **WKL**₀.*

Idea: Build a tree whose paths give perfect matchings.

Idea: Build a tree whose paths give perfect matchings.

$\langle a_0, a_1, \dots, a_n \rangle \in T$ iff $\{(0, a_0), (1, a_1), \dots, (n, a_n)\}$ is a matching.

Idea: Build a tree whose paths give perfect matchings.

$\langle a_0, a_1, \dots, a_n \rangle \in T$ iff $\{(0, a_0), (1, a_1), \dots, (n, a_n)\}$ is a matching.

Condition (A) guarantees the tree will be infinite.

Reversals

Steffens for locally finite graphs implies ACA_0 :



Reversals

Steffens for locally finite graphs implies ACA_0 :



Reversals

Steffens for locally finite graphs implies ACA_0 :




Reversals

Steffens for locally finite graphs implies ACA_0 :



Steffens for bounded graphs implies WKL_0 :




Reversals

Steffens for locally finite graphs implies ACA_0 :



Steffens for bounded graphs implies WKL_0 :




Reversals

Steffens for locally finite graphs implies ACA_0 :




Steffens for bounded graphs implies WKL_0 :




Reversals

Steffens for locally finite graphs implies ACA_0 :



Steffens for bounded graphs implies WKL_0 :




Reversals

Steffens for locally finite graphs implies ACA_0 :



Steffens for bounded graphs implies WKL_0 :




Graphs in general

Why is this messy?


Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?




Graphs in general

Why is this messy?



Graphs in general

Why is this messy?



The problem: To use a larger matching, you must abandon a smaller matching.

Independent matchings

Definition

A matching M is *independent* provided there is no **proper** M -augmenting path starting at a vertex unmatched by M .

A subgraph is *independent* provided it has a perfect matching, and all perfect matchings are independent.

Independent matchings

Definition

A matching M is *independent* provided there is no **proper** M -augmenting path starting at a vertex unmatched by M .

A subgraph is *independent* provided it has a perfect matching, and all perfect matchings are independent.

Lemma

Every graph has has a maximal independent matching.

Independent matchings

Definition

A matching M is *independent* provided there is no **proper** M -augmenting path starting at a vertex unmatched by M .

A subgraph is *independent* provided it has a perfect matching, and all perfect matchings are independent.

Lemma

Every graph has has a maximal independent matching.

Proof.

Zorn's lemma.



Independent matchings

Definition

A matching M is *independent* provided there is no **proper** M -augmenting path starting at a vertex unmatched by M .

A subgraph is *independent* provided it has a perfect matching, and all perfect matchings are independent.

Lemma

Every graph has has a maximal independent matching.

Proof.

Work in a countably coded β_2 -model; build an increasing sequence of independent matchings; argue that the union is maximal. □

Independent matchings

Definition

A matching M is *independent* provided there is no **proper** M -augmenting path starting at a vertex unmatched by M .

A subgraph is *independent* provided it has a perfect matching, and all perfect matchings are independent.

Lemma

Every graph has has a maximal independent matching.

Proof.

Work in a countably coded β_2 -model; build an increasing sequence of independent matchings; argue that the union is maximal. □

Note: this is a proof in Π_2^1 -CA.

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .
- ▶ If there is some v not matched, take a maximal independent matching M' of $G \setminus (V(M) \cup \{v\})$.

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .
- ▶ If there is some v not matched, take a maximal independent matching M' of $G \setminus (V(M) \cup \{v\})$.
- ▶ Use an M' -augmenting path starting at v to get a matching that includes M and v , and whose complement satisfies condition (A).

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .
- ▶ If there is some v not matched, take a maximal independent matching M' of $G \setminus (V(M) \cup \{v\})$.
- ▶ Use an M' -augmenting path starting at v to get a matching that includes M and v , and whose complement satisfies condition (A).
- ▶ Repeat.

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .
- ▶ If there is some v not matched, take a maximal independent matching M' of $G \setminus (V(M) \cup \{v\})$.
- ▶ Use an M' -augmenting path starting at v to get a matching that includes M and v , and whose complement satisfies condition (A).
- ▶ Repeat.

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .
- ▶ If there is some v not matched, take a maximal independent matching M' of $G \setminus (V(M) \cup \{v\})$.
- ▶ Use an M' -augmenting path starting at v to get a matching that includes M and v , and whose complement satisfies condition (A).
- ▶ Repeat.

Note: we potentially need the maximal independent subgraph lemma infinitely often,...

A proof of Steffens' theorem

Suppose G satisfies condition (A)

- ▶ Take a maximal independent matching M .
- ▶ If there is some v not matched, take a maximal independent matching M' of $G \setminus (V(M) \cup \{v\})$.
- ▶ Use an M' -augmenting path starting at v to get a matching that includes M and v , and whose complement satisfies condition (A).
- ▶ Repeat.

Note: we potentially need the maximal independent subgraph lemma infinitely often, but actually, exactly once.

A proof of maximality

Take a maximal independent subgraph H of G .

A proof of maximality

Take a maximal independent subgraph H of G .

Let N be the set of vertices not in H but adjacent only to vertices in H .

A proof of maximality

Take a maximal independent subgraph H of G .

Let N be the set of vertices not in H but adjacent only to vertices in H .

$G \setminus (H \cup N)$ satisfies condition (A), so by Steffens, has a perfect matching.

A proof of maximality

Take a maximal independent subgraph H of G .

Let N be the set of vertices not in H but adjacent only to vertices in H .

$G \setminus (H \cup N)$ satisfies condition (A), so by Steffens, has a perfect matching.

But the perfect matching would be independent in G , giving a larger independent matching. So any perfect matching of H is a maximal matching of G .

Reversals



Lemma

Given any tree T , there is a tree T' such that T has an infinite path iff T' has a perfect matching.

Reversals

Lemma

Given any tree T , there is a tree T' such that T has an infinite path iff T' has a perfect matching.





Proposition

Maximality implies Π_1^1 -CA₀. Steffens implies Σ_1^1 -AC₀.

Reversals

Lemma

Given any tree T , there is a tree T' such that T has an infinite path iff T' has a perfect matching.





Proposition

Maximality implies Π_1^1 -CA₀. Steffens implies Σ_1^1 -AC₀.

Reversals

Lemma

Given any tree T , there is a tree T' such that T has an infinite path iff T' has a perfect matching.



Proposition

Maximality implies Π_1^1 -CA₀. Steffens implies Σ_1^1 -AC₀.

Can we do better?


For any computable ordinal α , there is a computable graph G that satisfies condition (A), any perfect matching of which computes $\mathbf{0}^{(\alpha)}$.

Can we do better?

For any computable ordinal α , there is a computable graph G that satisfies condition (A), any perfect matching of which computes $\mathbf{0}^{(\alpha)}$.

This would be enough to prove ATR_0 , except we don't know how to prove G satisfies condition (A) without using $\Pi_1^1\text{-TI}_0$.
($\Sigma_1^1\text{-DC}_0$)

The current picture



The End

Thanks!